

Peru Conservation

Recorded Wildlife at Taricaya



Butterflies (Mariposas) in Taricaya Reserve, Madre de Dios

CLASS: Insecta

ORDER: Lepidoptera

1. Familia Nymphalidae

Subfamilia Apaturinae	
Doxocopa kallina	(Staudinger, 1886).
Doxocopa laure	(Drury, 1776).
Doxocopa lavinia	(Butler, 1886).
Doxocopa linda	(C. Felder & R. Felder, 1860).
Doxocopa pavon	(Latreille, 1809).
Subfamilia Nymphalinae	
Tribu Coeini	
Baetus aelius	(Stoll, 1780).
Baetus deucaliom	(C. Felder & R. Felder, 1860).
Baetus japetus	(Staudinger, 1885).
Colobura annulata	(Willmot, Constantino & J. Hall, 2001).
Colobura dirce	(Linnaeus, 1758).
Historis acheronta	(Fabricius, 1775).
Historis odius	(Fabricius, 1775).

Smyrna blomfilda	(Fabricius, 1781).
Tigridia acesa	(Linnaeus, 1758).
Tribu Kallimini	
Anartia jatrophae	(Linnaeus, 1763).
Junonia everate	(Cramer, 1779).
Junonia genoveva	(Cramer, 1780).
Metamorpha elissa	(Hübner, 1818).
Siproeta stelenes	(Linnaeus, 1758).
Tribu Melitaeini	
Eresia clio	(Linnaeus, 1758).
Eresia eunice	(Hübner, 1807).
Eresia nauplios	(Linnaeus, 1758).
Tegosa claudina	(Escholtz, 1821).
Tegosa fragilis	(H. W. Bates, 1864).
Tribu Nymphalini	
Hypanarthis lethe	(Fabricius, 1793).
Tribu Acraeini	
Actinote pellenea	(Hübner, 1821).
Subfamilia Charaxinae	
Tribu Preponini	
Agrias amydon	(Hewitson, 1854).
Agrias claudina	(Godart, 1824).

Archaeoprepona amphimacus	(Fabricius, 1775).
Archaeoprepona demophon	(Linnaeus, 1758).
Archaeoprepona meander	(Cramer, 1775).
Prepona dexamenus	(Hopffer, 1874).
Prepona laertes	(Hübner, 1811).
Prepona pheridamas	(Cramer, 1777).
Prepona pylene	(Hewitson, 1854).
Tribu Ananeini	
Consul fabius	(Cramer, 1776).
Fountainea eurypyle	(C. Felder & R. Felder, 1862).
Fountainea ryphea	(Cramer, 1775).
Memphis acidalia	(Hübner, 1819).
Memphis moruus	(Fabricius, 1775).
Memphis oenomais	(Boisduval, 1780).
Memphis phantes	(Hopffer, 1874).
Memphis philumena	(Doubleday, 1849).
Memphis polycarmes	(Fabricius, 1775).
Zaretis itys	(Cramer, 1777).
Subfamilia Morphinae	
Tribu Brassolini	
Bia actorion	(Linnaeus, 1763).

Brassolis sophorae	(Linnaeus, 1758).
Caligo eurilochus	(Cramer, 1775).
Caligo idomeneus	(Linnaeus, 1758).
Caligo illoneus	(Cramer, 1775).
Caligo teucer	(Linnaeus, 1758).
Catoblepia berecynthia	(Cramer, 1777).
Catoblepia soranus	(Westwood, 1851).
Catoblepia xanthicles	(Godman & Salvin, 1881).
Eryphanes automedon	(Cramer, 1775).
Narope nesope	(Hewitson, 1869).
Opotera aorsa	(Godart, 1824).
Opsiphanes cassiae	(Linnaeus, 1758).
Opsiphanes cassina	(C. Felder & R. Felder, 1862).
Opsiphanes invirae	(Hübner, 1808).
Opsiphanes quiteria	(Stoll, 1780).
Selenophanes cassiope	(Cramer, 1775).
Tribu Morphini	
Antirrhea hela	(C. Felder & R. Felder, 1862).
Caeris chorinaeus	(Fabricius, 1775).
Morpho deidamia	(Hübner, 1819).
Morpho helenor	(Cramer, 1776).

Morpho menelaus	(Linnaeus, 1758).
Morpoho achilles	(Linnaeus, 1758).
Subfamilia Satyrinae	
Cepheuptychia cephus	(J.C. Fabricius, 1775).
Caeruleuptychia lobelia	(Butler, 1870).
Chloroeuptychia catharina	(Staudinger, 1886).
Chloroeuptychia herseis	(Godart, 1824).
Chloroeuptychia hewitsoni	(Butler, 1867).
Chloroeuptychia sp.2	-
Cissia myneca	(Cramer, 1780).
Cissia palliada	(Butler, 1867).
Cissia proba	(Weymer, 1911).
Cissia sp.	-
Cissia terrestris	(Butler, 1867).
Cithaerias pireta	(Stoll, 1780).
Erichtodes antonina	(C. Felder & R. Felder, 1867).
Euptychia enyo	(Butler, 1818).
Euptychia sp1	-
Haetera Piera	(Linnaeus, 1758).
Harjesia blanda	(Moschler, 1964).
Hermeuptychia hermes	(Fabricius, 1775).

Magneptychia iris	(C. Felder & R. Felder, 1867).
Magneptychia libye	(Linnaeus, 1767).
Magneptychia modesta	(Butler, 1867).
Manataria hercyna	(Hübner, 1821).
Megeptychia antonoe	(Cramer, 1775).
Pareptychia binocula	(Butler, 1868).
Pareptychia hesionides	(Forster, 1964).
Pareptychia ocirrhoe	(Fabricius, 1776).
Pareptychia summandosa	(Gosse, 1880).
Paryphthimoides melobosis	(Capronnier, 1874).
Paryphthimoides melobosis sp. binalinea	(Capronnier, 1874).
Pierella hortona	(Hewitson, 1854).
Pierella hyalinus	(Gmelin, 1790).
Pierella lamia	(Sulzer, 1776).
Pierella lena	(Linnaeus, 1767).
Postaygetis penelea	(Cramer, 1777).
Pseudodebis marpesa	(Hewitson, 1862).
Pseudodebis valentina	(Linnaeus, 1763).
Splendeptychia aurigera	(Weymer, 1911).
Splendeptychia itonis	(Hewitson, 1862).
Taygetis angulosa	(Weymer, 1907).

Taygetis echo	(Cramer, 1775).
Taygetis imperator	New species discovered 2004
Taygetis laches	(Fabricius, 1793).
Taygetis larua	(C. Felder & R. Felder, 1867).
Taygetis mermeria	(Cramer, 1776).
Taygetis sylvia	(H. W. Bates, 1866).
Taygetis virgilia	(Cramer, 1776).
Taygetomorpha celia	(Cramer, 1779).
Ypthimoides mythra	(Weymer, 1911).
Ypthimoides sp.	-
Subfamilia Biblidinae	
Tribu Biblidini	
Biblis hyperia	(Cramer, 1779).
Callicore cinosura	(Doubleday, 1848).
Catoneohele numilia	(Cramer, 1775).
Catonephele acontius	(Linnaeus, 1771).
Catonephele antinoe	(Godart, 1824).
Catonephele salacia	(Hewitson, 1852).
Diaetria clymena	(Cramer, 1775).
Dynamine artemisia	(Fabricius, 1793).
Dynamine myrson	(Doubleday, 1849).

Dynamine racidula	(Hewitson, 1852).
Eunica alpais	(Godart, 1824).
Eunica eurota	(Cramer, 1775).
Eunica pusilla	(H. W. Bates, 1864).
Hamadryas amphinome	(Linnaeus, 1767).
Hamadryas arinome	(Lucas, 1853).
Hamadryas chloe	(Stoll, 1787).
Hamadryas feronia	(Linnaeus, 1758).
Hamadryas laodamia	(Cramer, 1777).
Myscelia capenas	(Hewitson, 1857).
Nessaea obrinus	(Linnaeus, 1758).
Nica flavilla	(Godart, 1824).
Panacea prola	(Doubleday, 1848).
Peria lamis	(Cramer, 1779).
Pyrrogyra amphiro	(Bates, 1865).
Pyrrogyra cramerii	(Aurivillus, 1882).
Pyrrogyra edocla	(Doubleday, 1848).
Pyrrogyra neaerea	(Linnaeus, 1758).
Pyrrogyra otolais	(H. W. Bates, 1864).
Temenis laothoe	(Cramer, 1777).
Vila azeca	(Doubleday, 1848).

Tribu Cyrestidini	
Marpesia berania	(Hewitson, 1852).
Marpesia chiron	(Fabricius, 1775).
Marpesia crethon	(Fabricius, 1776).
Marpesia egina	(H. W. Bates, 1865).
Marpesia furcula	(Fabricius, 1793).
Marpesia petreus	(Cramer, 1776).
Marpesia temístocles	(Fabricius, 1793).
Subfamilia Limenitidinae	
Adelpha amazona	(Austin & Jasinski, 1999).
Adelpha attica	(C. Felder & R. Felder, 1867).
Adelpha barnesia	(Schaus, 1902).
Adelpha boeotia	(C. Felder & R. Felder, 1867).
Adelpha capucinus	(Walch, 1775).
Adelpha cytherea	(Linnaeus, 1758).
Adelpha epione	(Godart, 1824).
Adelpha iphicleola	(H. W. Bates, 1862).
Adelpha jordani	(Frustorfer, 1912).
Adelpha malea	(C. Felder & R. Felder, 1861).
Adelpha melona	(Hewitson, 1847).
Adelpha mesentina	(Cramer, 1777).

Adelpha naxia	(C. Felder & R. Felder, 1867).
Adelpha paraena	(H. W. Bates, 1865).
Adelpha plesaure	(Hübner, 1823).
Adelpha thesprotia	(C. Felder & R. Felder, 1867).
Adelpha viola	(Frustorfer, 1913).
Adelpha ximena	(C. Felder & R. Felder, 1862).
Adeplha erotia	(Hewitson, 1847).
Subfamilia Heliconiinae	
Dione junio	(Cramer, 1779).
Dryas iulia	(Fabricius, 1775).
Euides aliphera	(Godart, 1819).
Euides isabella	(Stoll, 1781).
Helicobius melpómene	(Linnaeus, 1758).
Heliconis numata	(Cramer, 1780).
Heliconius erato	(Linnaeus, 1758).
Heliconius hecale	(Fabricius, 1776).
Heliconius sara	(Fabricius, 1793).
Heliconius wallacei	(Reakirt, 1866).
Laparus doris	(Linnaeus, 1761).
Philaethria dido	(Linnaeus, 1763).
Subfamilia Ithomiinae	

Ceratynia neso	(Hübner, 1806).
Episcada hymen	(Haensch, 1905).
Ithomia agnosia	(Hewitson, 1855).
McClugia sp.	-
Mechanitis lysimnia	(Fabricius, 1793).
Mechanitis mazaesus mazaesus	(Hewitson, 1860).
Mechanitis polimnia kayei	(R. M. Fox 1967).
Mechanitis polimnia	(Linnaeus, 1758).
Methona confusa	(Butler, 1873).
Napaeogenes pharo	(C. Felder & R. Felder, 1862).
Napeogenes inachia	(Hewitson, 1855).
Oleria onega	(Hewitson, 1852).
Oleria victorine	(Guerin-Meneville, 1844).
Pteronymia forsteri	(Bauman, 1985).
Scada reckia	(Hübner, 1808).
Thyridia psidii	(Linnaeus, 1758).
Tithorea harmonia	(Cramer, 1777).

2. Familia Pieridae

Subfamilia Pierinae	
Tribu Coliadinae	
Anteos menipe	(Hübner, 1818).

Aphrissa statira	(Cramer, 1777).
Eurema albula	(Cramer, 1775).
Eurema lirina	(H. W. Bates, 1861).
Eurema sp1	-
Eurema sp2	-
Eurema sp3	-
Eurema sp4	-
Phoebis argante	(Fabricius, 1775).
Phoebis philea	(Linnaeus, 1763).
Phoebis sennae	(Linnaeus, 1758).
Rhabdodryas trite	(Linnaeus, 1758).
Tribu Pierinae	
Ascia monuste	(Linnaeus, 1777).
Ganyra phaloe	(Godart, 1819).
Glutophrissa drusilla	(Cramer, 1777).
Itaballia demophile	(Linnaeus, 1763).
Itaballia pandosia	(Hewitson, 1853).
Melete lycimnia	(Cramer, 1777).
Perrhybris pamela	(Stoll, 1780).
Pieriballia viardi	(Boisduval, 1836).

Pyristidia nise	(Cramer, 1775).
Tribu Dismorphiinae	
Enantia lina	(Herbst, 1792).
Enantia melite	(Linnaeus, 1763).
Pseudopieris nehemia	(Boisduval, 1836).

3. Familia Riodinidae

Subfamilia Riodininae	
Argyrogrammana stilbe	(J.B. Godart, 1824).
Adelotypa sejuncta	(Stichel, 1910).
Adelotypa huebneri	(A.G. Butler, 1867).
Alesa lipara	(H. W. Bates, 1867).
Anteros acheus	(C. Stoll, 1781).
Anteros formosus	(P. Cramer, 1777).
Anteros renaldus	(C. Stoll, 1790).
Baeotis staudingeri	(B.L. D'Abrera, 1994).
Caria sp.	-
Cariomothis erytromelas	(Sepp, 1841).
Cyrenia martia	(Westwood, 1851).
Detritivora off. cleonus	(Stoll, 1781).
Emesis castigata	(Stichel, 1810).

Emesis mandana	(Cramer, 1780).
Euselasia violetta	(H. W. Bates, 1868).
Euselasia calligramma	(H.W. Bates, 1868).
Euselasia ignitus	(H.F.E.J. Stichel, 1924).
Euselasia serapis	(H.F.E.J. Stichel, 1919).
Euselasia euphaes fulviplaga	(W.C. Hewitson, [1855])).
Euselasia eustola eustola	(H.F.E.J. Stichel, 1919).
Lasaia agesialas	(Latreille, 1809).
Livendula aminias	New species discovered 2004
Livendula huebneri	New species discovered 2004
Livendula balista	New species discovered 2004
Livendula sp. 1	-
Mesene epaphus	(C. Stoll, 1780).
Mesene leucophrys	(H.W. Bates, 1868).
Mesene silaris	(Godman & Salvin, 1878).
Mesosemia naiadella	(Stichel, 1909).
Mesosemia thymetus	(Cramer, 1777).
Mesosemia judicialis	(Butler, 1874).
Mesosemia maeotis	(Hewitson, 1859).
Mesosemia philocles	(Linnaeus, 1758).

Mesosemia sp1	-
Mesosemia sp3	-
Mesosemia thetys	(Godman & Salvin).
Metacharis lucius	(Fabricius, 1793).
Metacharis regalis	(Butler, 1867).
Napaea beltiana	(H.W. Bates, 1867).
Napaea heteroea	(H.W. Bates, 1867).
Nymphidium acherois	(Boisduval, 1836).
Nymphidium ascolia	(Hewitson, 1853).
Nymphidium aurum	(C.J. Callaghan, 1985).
Nymphidium azanoides	(A.G. Butler, 1867).
Nymphidium baeotia	(Hewitson, 1853).
Nymphidium caricae	(Linnaeus, 1758).
Nymphidium hesperinum	(H.F.E.J. Stichel, 1911).
Nymphidium fulminans	(H.W. Bates, 1868).
Nymphidium leucosia medusa	(H. Druce, 1904).
Nymphidium ninias	(Hewitson, 1865).
Nymphidium sp. 1.	-
Nothome erota	(Cramer, 1780).
Parcella amarynthina	(C. Felder & R. Felder, 1865).

Rethus arcus	(Linnaeus, 1763).
Rethus periander	(Cramer, 1777).
Sarota spicata	(Staudinger, 1888).
Setabis sp.	(Westwood, 1851).
Synargis calyce	(C. Felder & R. Felder, 1862).

4. Familia Lycaenidae

Subfamilia Teclinae	
Celmia celmus	(Cramer, 1775).
Strymon ziba	(Hewitson, 1868).

5. Familia Hesperidae

Astrartes anaphus	(P. Cramer, 1777).
Astrartes sp.	-
Astrasptes creteus	(P. Cramer, 1780).
Autochton sp. 1	-
Gorgopas trochillus	(C.H. Hopffer, 1874).
Nascus sp. 1	
Nascus phocus	(P. Cramer, 1777).
Nisionades sp.1	-
Phocides metrodurus	(E.L. Bell, 1932).
Urbanus teleus	(J. Hübner, 1821).

6. Familia Papilionidae

Heraclides astyalus	(Godart, 1819).
Mimoides ariarathes	(Esper, 1788).
Mimoides pausianas	(Hewitson, 1852).
Parides lysander	(Cramer, 1775).
Parides lysander brissonius	(Hübner, 1819).
Parides neophilus	(Geyer, 1837).
Parides neophilus olivencius	(Geyer, 1837).
Parides vertumnus astorius	(J. Zicán, 1940).
Parides vertumnus	(Cramer, 1779).
Protesilaus glaucolaus	(H. W. Bates, 1864).
Protographium agesilaus	(Guerin-Meneville & Percheron, 1835).

Total 281 species